

**B.Sc. PHYSICS
FIFTH SEMESTER
CLASSICAL DYNAMICS
BSP-503A [SPECIAL REPEAT]
[USE OMR FOR OBJECTIVE PART]**

**SET
A**

Duration: 3 hrs.

Full Marks: 70

Time: 30 min.

(Objective)

Marks: 20

Choose the correct answer from the following: 1X20=20

- How many independent variables are there in the Hamiltonian $H=H(q_k, p_k, t)$, where $k=1,2,\dots,n$?
 - n
 - n+1
 - 2 n
 - 2 n+1
- For a parabolic path of planets, the eccentricity parameter ϵ will
 - 0
 - 1
 - >1
 - <1
- The Lagrangian of a system is $L = m \dot{q}^2$, its Hamiltonian will be
 - $\frac{p^2}{2m}$
 - $\frac{2p^2}{m}$
 - $\frac{p^2}{4m}$
 - $\frac{p^2}{m}$
- In a conservative system, the Lagrangian equations of motions will be
 - $\frac{d}{dt} \left(\frac{\partial T}{\partial \dot{q}_k} \right) = \frac{\partial T}{\partial q_k}$
 - $\frac{d}{dt} \left(\frac{\partial L}{\partial \dot{q}_k} \right) = \frac{\partial L}{\partial q_k}$
 - $\frac{d}{dt} \left(\frac{\partial T}{\partial \dot{q}_k} \right) = \frac{\partial T}{\partial q_k}$
 - $\frac{d}{dt} \left(\frac{\partial L}{\partial \dot{q}_k} \right) = \frac{\partial L}{\partial q_k}$
- A particle moves in a plane. It's kinetic energy will be
 - $T = \frac{1}{2} m (\dot{r}^2 + r \dot{\theta}^2)$
 - $T = \frac{1}{2} m (\dot{r}^2 + r^2 \dot{\theta})$
 - $T = \frac{1}{2} m (\dot{r}^2 + r^2 \dot{\theta}^2)$
 - $T = \frac{1}{2} m (\dot{r}^2 + \dot{\theta}^2)$
- A central force acting on a particle is given by $F \propto -1/r^2$, potential of the system will be
 - $V \propto -\frac{1}{r}$
 - $V \propto \frac{1}{r}$
 - $V \propto -r$
 - $V \propto r$
- The potential energy of a system is given by $V = mgz$, then the force acting on the particle will be
 - $\vec{F} = mg\hat{z}$
 - $\vec{F} = -mg\hat{z}$
 - $\vec{F} = m g \hat{z}$
 - $\vec{F} = -m g \hat{z}$
- Which one of the following is correct for the Hamiltonian equations of motion
 - $\dot{p}_k = -\frac{\partial H}{\partial q_k}$
 - $\dot{q}_k = -\frac{\partial H}{\partial p_k}$
 - $\dot{q}_k = \frac{\partial H}{\partial p_k}$
 - $\dot{p}_k = \frac{\partial H}{\partial q_k}$

9. Consider two frames S_1 and S_2 , where the later one is moving with a relativistic velocity v along a particular direction. Statements: (i) two events are simultaneous in S_1 frame, then (ii) these two events are not simultaneous in S_2 frame
- Both statements are true
 - Statements (i) is true & (ii) is false
 - Statements (i) is false & (ii) is true
 - Both statements are false
10. A 500m long train moving with a speed $0.8c$, its moving length will be
- 200m
 - 250m
 - 300m
 - 500m
11. The reduced mass of positronium atom will be
- $\frac{m_e}{2}$
 - $\frac{3m_e}{2}$
 - m_e
 - $2m_e$
12. Two photons approaching each other. Their relative velocity will be
- $c/2$
 - c
 - $2c$
 - $c/3$
13. The interval between two events will be light-like in Minkowski space-time provided
- $ds^2 > 0$
 - $ds^2 < 0$
 - $ds^2 = 0$
 - $ds^2 \leq 0$
14. A moving clock appears
- slow
 - fast
 - Remain same
 - Sometimes fast and sometimes slow
15. The aerial velocity is defined by
- $r^2 \dot{\theta}$
 - $\frac{1}{2} r^2 \dot{\theta}$
 - $\frac{1}{2} r^2 \dot{\theta}^2$
 - $r^2 \dot{\theta}^2$
16. The Hamiltonian of a system will defined by
- $H = \sum p_k \dot{q}_k + L(q_k, \dot{q}_k, t)$
 - $H = -\sum p_k \dot{q}_k - L(q_k, \dot{q}_k, t)$
 - $H = L(q_k, \dot{q}_k, t) - \sum p_k \dot{q}_k$
 - $H = \sum p_k \dot{q}_k - L(q_k, \dot{q}_k, t)$
17. The relativistic energy-momentum relation will be
- $E = \pm \sqrt{p^2 c^2 + m^2 c^4}$
 - $E = \pm \sqrt{p^2 c^4 + m^2 c^2}$
 - $E = \pm \sqrt{p^2 c^4 + m c^2}$
 - $E = \pm \sqrt{p^2 c^2 + m^2 c^4}$
18. The total energy of a two-body problem will be
- $\frac{1}{2} \mu r^2 + \frac{1}{2} \frac{J^2}{\mu r^2} + V$
 - $\frac{1}{2} \mu r^2 + \frac{1}{2} \frac{J^2}{\mu r^4} + V$
 - $\frac{1}{2} \mu r^2 + \frac{J^2}{\mu r^2} + V$
 - $\frac{1}{2} \mu r^2 + \frac{1}{2} \frac{J^2}{\mu r^2} + V$
19. The angular momentum of two-body problem will be
- $\dot{\theta} = \frac{J}{\mu r^2}$
 - $J = \frac{\dot{\theta}}{\mu r^2}$
 - $J = \mu r^2 \dot{\theta}^2$
 - $\dot{\theta} = \mu r^2 / J$

20. The total energy of planets revolving around the sun in a circular paths will be

a. $E = -\frac{\mu k^2}{J^2}$

b. $E = -\frac{\mu k^2}{2 J^2}$

c. $E = -\frac{\mu J^2}{k^2}$

d. $E = -\frac{\mu J^2}{2 k^2}$

(Descriptive)

Time : 2 hrs. 30 mins.

Marks : 50

[Answer question no.1 & any four (4) from the rest]

- | | | |
|-------|--|--------------|
| 1. a. | Show that path of planets under a central force motion is conic section. | 8+2=10 |
| b. | State the conditions of parabolic and circular paths of the planets. | |
| 2. a. | A particles moves in plane under a central potential $V = -\frac{k}{r}$. Find the Lagrangian of the system. | 2+6+2
=10 |
| b. | Obtain the Lagrange's equations of motions. | |
| c. | Find the Hamiltonian of the system. | |
| 3. a. | Construct the Lagrangian of a simple pendulum. | 3+5+2
=10 |
| b. | Find the Lagrange's equations of motions of a simple pendulum. | |
| c. | For a small oscillation of a simple pendulum, find its time period. | |
| 4. a. | State Kepler's laws of planetary motions. | 4+3+3
=10 |
| b. | Prove the laws of period of Kepler's law. | |
| c. | If the earth suddenly shrinks to its radius by 25%, calculate the new time period. | |

5. a. Derive the components of acceleration of a particle moves in a plane. 5+2+3
=10
 b. Show that the angular momentum in a two-body system is conserved.
 c. Find an expression of effective potential energy of this two-body system.
6. a. Define Hamiltonian of a system. 2+4+4
=10
 b. Show that the Hamiltonian of a system is $H = \sum p_k \dot{q}_k - L$.
 c. Derive the Hamiltonian equations of motions.
7. a. Using Lorentz transformation relations derive the velocity addition theorem. 4+2+4
=10
 b. Show that the speed of light in vacuum is the upper limit of speed of any object.
 c. If the time interval measured by a moving observer is 3 hrs when moving with a relativistic velocity $0.8c$, find its proper time.
8. a. Using Lorentz transformation relations show that $ds^2 = -c^2 dt^2 + dx^2 + dy^2 + dz^2$ is invariant. 4+3+3
=10
 b. The total energy of a relativistic particle is 2 times its rest mass energy. What will be its speed?
 c. Derive an expression of time dilation in special relativity theory.

== *** ==